

Delivery Management for HIV Positive Women

MOTHER

ANTEPARTUM

Begin mulidrug regimen of three drugs at 12 weeks gestation (or immediately if ill) or continue if currently being treated.

INTRAPARTUM

Continue PO HIV meds as able and add Zidovudine (AZT)

Loading Dose:

2mg/kg = _____ mg
infused over one hour

Second Dose:

1mg/kg = _____ mg
per hour until delivery

PREPARATION

Zidovudine (AZT)
1000 mg/250 ml D5W

Final concentration:
4mg/ml

QUESTIONS?

Please refer all HIV pregnancy care to Northern Nevada HOPES.

- Contact our RNs for a **detailed consultation** on **procedures** and **discharge planning**: 775-997-7537
- Contact our Women/Infant Case Manager for all other general inquires: 775-223-9135
- Contact our Case Managers if you are having trouble reaching our specialists: 775-348-2893

INFANT

- Draw HIV RNA by PCR at birth
- Draw HIV Elisa at birth

FOR PO INFANT

Zidovudine (AZT)

4mg/kg = _____ mg
PO q 12 hours for six weeks

- Start within 6 – 12 hours of delivery

FOR NPO INFANT

Zidovudine (AZT)

3mg/kg = _____ mg
IV q 12 hours.

Infused over 30 – 60 minutes

- Start within 6 – 12 hours of delivery

If infant < 35 wks. call
for consultation

