

# Delivery Management for HIV Positive Women

## MOTHER

### ANTEPARTUM

Begin mulidrug regimen of three drugs at 12 weeks gestation (or immediately if ill) or continue if currently being treated.

### INTRAPARTUM

Continue PO HIV meds as able and add Zidovudine (AZT)

#### **Loading Dose:**

2mg/kg = \_\_\_\_\_ mg  
infused over one hour

#### **Second Dose:**

1mg/kg = \_\_\_\_\_ mg  
per hour until delivery

### PREPARATION

Zidovudine (AZT)  
1000 mg/250 ml D5W

Final concentration:  
4mg/ml

## QUESTIONS?

Please refer all HIV pregnancy care to Northern Nevada HOPES (775-786-4673) and ask to speak with a Case Management RN for a detailed consultation.

## INFANT

- Draw HIV RNA by PCR at birth
- Draw HIV Elisa at birth

### FOR PO INFANT

Zidovudine (AZT)

4mg/kg = \_\_\_\_\_ mg  
PO q 12 hours for six weeks

- Start within 6 – 12 hours of delivery

### FOR NPO INFANT

Zidovudine (AZT)

3mg/kg = \_\_\_\_\_ mg  
IV q 12 hours.

Infused over 30 – 60 minutes

- Start within 6 – 12 hours of delivery

If infant < 35 wks. call  
for consultation

