

# Delivery Management for HIV Positive Women

## MOTHER

### ANTEPARTUM

Begin mulidrug regimen of three drugs at 12 weeks gestation (or immediately if ill) or continue if currently being treated.

### INTRAPARTUM

Continue PO HIV meds as able and add Zidovudine (AZT)

#### Loading Dose:

2mg/kg = \_\_\_\_\_ mg  
infused over one hour

#### Second Dose:

1mg/kg = \_\_\_\_\_ mg  
per hour until delivery

### PREPARATION

Zidovudine (AZT)  
1000 mg/250 ml D5W

Final concentration:  
4mg/ml



## INFANT

- Draw HIV RNA by PCR at birth
- Draw HIV Elisa at birth

### FOR PO INFANT

Zidovudine (AZT)

4mg/kg = \_\_\_\_\_ mg  
PO q 12 hours for six weeks

- Start within 6 – 12 hours of delivery

### FOR NPO INFANT

Zidovudine (AZT)

3mg/kg = \_\_\_\_\_ mg  
IV q 12 hours.

Infused over 30 – 60 minutes

- Start within 6 – 12 hours of delivery

If infant < 35 wks. call  
for consultation

## QUESTIONS?

Please refer all HIV pregnancy care to Northern Nevada HOPES: (775) 786-4673.

- Contact our RNs for a **detailed consultation** on **procedures** and **discharge planning**: (775) 786-4673 ext 1169.
- Contact our Case Managers if you are having trouble reaching our specialists: (775) 348-2893.